

**DETERMINATION OF AFLATOXIN M1
CONCENTRATIONS IN FULL-FAT COW'S UHT MILK
SOLD FOR CONSUMPTION IN NAJHRAN-SAUDI
REGARDING ITS PUBLIC HEALTH SIGNIFICANCE***

*Medhat I.M. Abdallah¹; Mohamed S. Bazalou²
and Mohamed Z. Al-Julaifi³*

***This study was supported by Deanship of Scientific Research Najran University
(Research Code No.: NU 67/11)**

- ¹ Associated Professor of Microbiology at Medical Laboratory Technology (MLT) Program, Department of Applied Medical Science, Community College (NCC), Najran University (NU), Najran, Kingdom of Saudi Arabia.
- ² Senior of Veterinary Specialist and food Inspector at Damietta Sea' Port Lab; Researcher (Ph.D.) in Biochemistry. Also, A Staff Member in Toxins, Hormones and Drugs Residues Lab at Ministry of Agriculture, Riyadh, Kingdom of Saudi Arabia.
- ³ Professor of Science, Molecular, Biotechnology and National Director of Project (UTF/SAU/035/SAU) Improving Animal Disease and Vaccine Production in the Kingdom of Saudi Arabia between Saudi Arabia and FAO. Also The General Director of Veterinary Laboratories Department at Ministry of Agriculture, Riyadh, Kingdom of Saudi Arabia.

ABSTRACT

This study aimed to evaluate the concentrations of aflatoxin M1 in full fat, cow's UHT milk solid in Najran–Saudi Arabia with regard to its public health significance. 96 samples of different brands full fat, cow's UHT milk were randomly punched from different supermarkets at Najran city during the period of September 2011 to January 2012. The samples were examined for AFM1 using the competitive enzyme-linked immunosorbent assay (ELISA), AFM1 residues were detected in 79 samples (82.30% of total). The minimum concentration was 0.01, the maximum concentration was 0.19 and the mean value was $0.058 \pm 0.0053 \mu\text{g/L}^{-1}$. Data also indicated that AFM1 residues concentrations detected in all the positive samples were below the tolerated level of AFM1. So it could be concluded that contamination of AFM1 in dairy products marketed in Najran city does not appear to be serious public health problem at the moment.

¹ Senior Researcher of Food Inspection at Damietta Sea' Port Lab, Reference Laboratory for Safety Analysis of Food from Animal Origin, Animal Health Research Institute (AHRI), Agricultural Research Center (ARC), Giza, Egypt.

INTRODUCTION

Good health starts with good nutrition and good nutrition can protect against diseases later in life. Liquid milk is a common health drink consumed by people of all age groups especially children. A large population in our countries depends on milk from local suppliers. Milk is a product of biological evolution, its role in human nutrition is well known and its biochemical complex which appears to be the only material to function solely as a source of food. The complements of proteins in milk are ideal in quality and balance to satisfy human amino acid requirements. Confirmation of this nutritive image is the widespread use of milk and milk products as a part of the daily diet of peoples in the highly developed countries. As a consequence, such societies enjoy almost complete freedom from nutritional disease among infants, children, young and adults. In contrast, the underdeveloped areas of the world have a primitive or nonexistent milk supply and have numerous inhabitants suffering from nutrient deficiencies, especially infants and children (Cashman 2002 and Hoppe et al. 2006).

Human health is highly attractive world, so food safety remains a major challenge to food producers and to legislators endeavoring to adequate consumer protection. Both man and animals live under a certain degree of "biological hazard" from natural toxicants that occur in food and foodstuffs (Allam et al. 1999-a; 1999-b & 2002 and Abdelhamid et al. 2002).

Naturally occurring toxins such as mycotoxins pose intense challenges to food safety widespread in many countries, especially in tropical and subtropical regions where temperature and humidity conditions are optimum for growth of moulds and toxins production, so they are found in a wide variety of agricultural products (such as corn, wheat, soybean, barley and rice) (FAO (1991), and animal foods as well as meat products, milk products including ultra-high treated (UHT) milk and as a result of carry-over from contaminated animal feed (Blunden et al. 1991; Moustafa 1994; Adams & Moss 1995; Orriss 1997; Sayed et al. 2000; Takahashi-Ando et al. 2004; Cavaliere et al. 2006 and Trucksess et al. 2006). Mycotoxins contamination of food and feeds remains a worldwide problem, the United Nation Food and Agriculture Organization (FAO) has estimated that up to 25% of the world's food crops are significantly contaminated with mycotoxins (Jelinek et al. 1989; Smith et al. 1994 and WHO 1999). Mycotoxins are unavoidable food contaminants even when good agricultural practices are applied.

Crop transfers through international trade have made aflatoxins contaminated food a worldwide problem (Sherif 2003).

Nowadays the main mycotoxins of interest are aflatoxins, ochratoxins, trichothecenes, zearalenone, fumonisins, ergot alkaloids and deoxynivalenol (Hussein & Brasel 2001; Bhat & Vasanthi 2003 and Cleveland et al. 2003). Aflatoxins (AFs) are a group of polyketide-derived furanocoumarins, with at least 16 structurally related toxins that have been characterized, these toxins are produced by a number of different *Aspergillus* species (CAST 1989; FAO 1991; Abdelhamid et al. 1993; Goto et al. 1996; Klich et al. 2000; Ito et al. 2001 and Peterson et al. 2001). However, in the agriculture commodities, they are primarily produced by *Aspergillus flavus*; *Aspergillus parasiticus* and *Aspergillus nomius* (Moss 1998; Galvano et al. 2001; Creppy 2002 and Oliveira & Ferraz 2007). There are four major categories of AFs (AFB1, AFB2, AFG1 & AFG2) all of which occur naturally (Anonymous 1998) and according to the research by joint (FAO/WHO 1996) AFB1 is usually predominant of mould producing compounds and the most toxic of the four categories (Bhat & Miller 1991 and Brera et al. 1998). Meanwhile, AFB1 and AFB2 have assumed the biosynthetic precursor of the other significant members of aflatoxins family (Dutton et al. 1985), M1 and M2 are 4-hydroxy derivatives of AFB1 and AFB2 respectively (Chifiriuc et al. 2010). Several research workers reported that there is a linear relationship between the amount of AFM1 in milk and AFB1 in feed which is consumed by dairy cattle (Bakirci 2001). The half-time of AFM1 in milk is relatively short, if the intake of AFB1 ceases, AFM1 levels in milk are reduced by about 50% within 25-36 hours (FAO/WHO 1996) while, decrease to an untraceable level within 72 hours after removal of contaminated source (Rahimi & Karim 2008). In fact, AFM1 has a high affinity for casein (Brackett & Marth 1982; Galvano et al. 1998; Baskaya et al. 2006 and Barbiroli et al. 2007) and can be detected in milk through 12-24 hours as a direct result of ingestion of food contaminated with aflatoxin B1 by many mammals including dairy cattle and humans (Stoloff 1977 and Ghazani 2009). The conversion rate of aflatoxin B1 to aflatoxin M1 present in milk ranges between 0.5% and 5%, yet values as high as 6% have been reported by Pipet (1998).

AFM1 recognized as milk toxin exhibits, carcinogenic (IARC 2002), genotoxic (Lafont et al. 1989) and cytotoxic effects (Neal et al. 1998). AFB1 and AFM1 have been classified by the International Agency for Research on Cancer as human carcinogens class 1A (carcinogen) and 2B (probable carcinogen) respectively (IARC 2002),

because of their potential risks and also to minimize their hazard, the World Health Organization recommends the reduction in its consumption to a minimum, because there is still not enough information available to establish a tolerable exposure level (WHO 2002). Fortunately, AFM1 has been found to be about 10 times less carcinogenic than AFB1 (Lafont et al. 1989). Researchers from US Division of Epidemiology and Surveillance have articulated that the ingestion of mycotoxin-contaminated animal-based food products could pose a concern to public health (Hollinger & Ekperigin 1999).

AFM1 is relatively stable in raw and processed milk products, pasteurization, sterilization and ultra-high-temperature (UHT) treatment or processing result in negligible destruction of AFM1 (Galvano et al. 1996; Yaroglu et al. 2005 and Unusan 2006). It has been reported that AFM1 was a resistant to thermal inactivation during food processing for procedures such as pasteurization and autoclaving (Bakirci 2001) and Park (2002). This toxin was found in foods in spite of disappearing fungal organisms (Macdonald & Castle 1996).

To secure safety of foodstuffs, regular monitoring of mycotoxins is necessary and for this purpose the following work is to provide a qualitative overview about aflatoxin M1 residues in marked UHT milk at Najran city, Saudi Arabia.

MATERIAL AND METHODS

Sampling:

A total of 96 samples of different brands full-fat cow's UHT milk, were randomly purchased from different supermarkets in various districts of Najran city in the period from September 2011 to January 2012. The samples were labeled, collected and transported to the laboratory in an insulated container at about 4°C and analyzed before their expiration dates exceeded.

Methods:

The samples were examined for AFM1 using the competitive enzyme-linked immunosorbent assay (ELISA), (RIDASCREEN Aflatoxin M1, R-Biopharm) test kits and the procedures were as described by Biopharm and Gmb (1999). This method is quick, reliable and cost effective for estimation of AFM1 and has been included in the official collection of test procedures by the German Federal Board of Health. All UHT milk samples were prepared and defatted using the method outlined in the ELISA kits, as briefly described. The mean lower detection limit of RIDASCREEN® AFM1 test is 0.01 µg/L⁻¹ for milk.

Statistical analysis:

Analysis of data was performed according statistical programmed of **SPSS (1997)**.

RESULTS AND DISCUSSION

Data outlined in table, 1 and 2 showed the analysis of 96 samples of commercial UHT whole milk, AFM1 residues were detected in 79 samples (82.30%) of total in different levels as showed in table, (2) with the mean value $0.058 \pm 0.0053 \mu\text{g/L}^{-1}$. Whereas, all the positive samples were below the limit permitted by **SASO (1996)**. Moreover, 17 samples (17.70%) total of UHT whole milk samples were not contain AFM1 as recorded in table, (1).

Table, (1): Occurrence and frequency distribution of examined “UHT milk” samples based on their AFM1 concentration (n=96):

Full-fat cow's UHT milk examined samples	Not detected samples	Positive samples	Frequency distribution of examined samples according to their AFM1 concentrations ($\mu\text{g/L}^{-1}$)*		Min.	Max.	Mean \pm SE
			< 0.20	> 0.20			
Number	17.00	79.00	79.00	00.00	0.01	0.19	0.058 \pm 0.0053
Percentage	17.70%	82.30%	82.30%	00.00%			

*According to SASO (1996)

Table, (2): Occurrence and frequency distribution of positive examined “UHT milk” samples based on their AFM1 concentration (n=79):

Range of AFM1 ($\mu\text{g/L}^{-1}$)	No. of positive samples	Percentage*
0.01 - 0.04	24	30.38%
0.05 - 0.09	30	37.97%
0.10 - 0.14	19	24.05%
0.15 - 0.19	06	07.60%
Detected	79	100.00%

*According to $0.2 \mu\text{g/L}^{-1}$ limit; N0. 79 x < 0.20 (100%)

A number of countries particularly developing ones, have established acceptance levels for AFM1, from 0.05 ppb in most European countries to 0.5 ppb in the United States, (**Van Egmond 1989**). (**Chen & Gao 1993** and **Van Egmond 1995**) have reported that in case of AFM1 in liquid milk, regulatory limits throughout the world are influenced by economic considerations, degree of development and may vary from one country to another, the Saudi Arabia Standard Organization has laid down the limit for total aflatoxins in liquid milk and its products (except dried milk) as $0.2 \mu\text{g/L}^{-1}$, and for infants and children foods as $0.05 \mu\text{g/kg}^{-1}$, (**SASO 1996** and **Gulf Standard 1997**).

As far as we knew, no available information dealing with the occurrence of AFM1 in UHT milk in Saudi Arabia. Elsewhere, nearly similar findings were reported by **Martins & Martins (2000)**; **Roussi et al. (2002)** and **López et al. (2003)**. While, higher prevalence was revealed by **Kamkar (2008)**; **Tekinşen & Eken (2008)**; **Gundinc & Filazi (2009)**; **Atasever et al. (2010)** and **German et al. (2010)**. Moreover, lower findings were reported by **Unusan (2006)** and **Fallah (2010)**.

In the light of the above data, it could be noticed that the obtained positive results of this study were below the tolerated level of AFM1 in UHT milk collected from Najran city, Saudi Arabia, that is may be due to locality of Najran city in the south part of Saudi Arabia with a hot climatic to some extent, mostly all over the year. Factors include environmental temperature, humidity and moisture content of the feed as well as pH, affecting AFM1 production by the mold. Consequently, fungi present in haystacks may not easily produce high levels of toxins during storage conditions and following the consumption of contaminated feed with AFB1, conversion of AFB1 to AFM1 which takes place in the liver may not leads to elevated levels of AFM1 in milk (**Sassahara et al. 2005** and **Prandini et al. 2009**).

Regarding the public health significance, mycotoxins attract worldwide attention because of the significant associated with their impact on human health (**CTA 1997**). Mycotoxins in small amount contamination of feed and food is a current problem, however their continuous intake even in microdoses can result in their accumulation in the human bodies, during a long term consumption which can cause a variety of ill human effects ranging from allergic responses to immunosuppressive, mutagenic, teratogenic and carcinogenic effects especially on the liver, probably underestimates the effect of mycotoxins as a cause of human mortality (**Varman & Evans 1991**; **El-Shinawy et**

al. 1994; Peraica et al. 1999; Pitt 2000; Kocabas & Sekerel 2003; Kovacs 2004; Baskaya et al. 2006 and Chifiriuc et al. 2010). Aflatoxin B₁ and aflatoxin M₁ are known as hepatoxins and hepatocarcinogens. Their harmful effects in human, especially infants are vital (Qian et al. 1984 and Chu 1991). In developed countries, aflatoxin contamination rarely occurs at levels that cause notable aflatoxicosis in humans. Epidemiological studies done in Asia and Africa that have demonstrated a positive association between dietary aflatoxins and liver cell cancer (FAO/WHO 1996; Haggag et al. 2001; Bhat & Vasanthi 2003 and Sherif 2003). Severe aflatoxicosis, a pathological condition caused by aflatoxin intake is produced when moderate to high levels of aflatoxins are consumed, acute episodes of disease ensue and may include hemorrhage, vomiting, abdominal pain, alteration in digestion, absorption and/or metabolism of nutrients, stunted growth, pulmonary edema, acute liver damage, convulsions, coma, and possibly death with cerebral edema and fatty involvement of the liver, kidneys and heart. Chronic aflatoxicosis can result from ingestion of low to moderate levels of aflatoxins (USFDA 2000). The importance of aflatoxin is emphasized by a recent outbreak of aflatoxicosis in Kenya in April 2004 in which 125 people died (Lewis et al. 2005).

Synergistic effects may observe as a result from interaction of aflatoxins with hepatitis B virus in the etiology of liver cancer and could interact with HIV/AIDS (Montesano et al. 1997 and FAO 1997-b & 2001).

CONCLUSION AND RECOMMENDATION

The results of this study indicate that occurrence of aflatoxin M₁ concentrations in UHT milk samples sold in Najran region were slight, probably because of slight to moderate contamination feedstuff consumed by the milking cows. It can be concluded that contamination of aflatoxin M₁ in dairy products marketed in Najran city does not appear to be a serious public health problem at the moment, although the presence of this toxin in most sample analyzed. However, frequent analytical surveillance by food control agencies is highly recommended to control the incidence of mycotoxin contamination in Saudi Arabia, especially in dairy products. Implementing a food control system, such as the Hazard Analysis and Critical Control Point (HACCP) system in the food industries (FAO 2001), suggest an efficient means for limiting mycotoxin contamination in the Saudi's food supply.

The most effective way of controlling aflatoxin M1 in food supply is to reduce contamination of raw material and supplementary feedstuffs for dairy cattle with aflatoxin B1. Specific regulation exist in many countries (FAO/WHO 1996), and practical programs are being developed as the Codex Committee on Food Additives and Contaminants has developed, a code of practice for reducing aflatoxin B1 in raw materials (Van Egmond et al. 1997). Reduction can be achieved by good manufacturing practices and good storage practices.

ACKNOWLEDGEMENT

The authors wish to express their thanks and appreciations to **Dr. Abdel Monem Ghanim**, and **Dr. Mohamed W. Nofal**, our colleagues in the NCC at Najran University for their valuable guidance and assistance in completing the statistical part of this study. Also, deepest thank to **Dr. Ahmed Maher Nasr** for his assistance.

REFERENCES

- Abdelhamid, A.M; T.M. Dorra; S.E. Mansy and A.E. Sallam (1993):** Effect of raising dietary protein, amino acids and/or energy levels as an attempt to alleviate severity of the chronic aflatoxicosis by broiler I. On the performance and Toxicological Symptom., J. Egypt. Ger. Soc. 7001., 12: 277.
- Abdelhamid, A.M.; A.E. Sallam; G.A. Abd Allah and S.H. El-Samra (2002):** Effect of feeding male rats on aflatoxic diets without or with medicinal herbs (thyme, safflower, ginger, black cumin and/or garlic)., Proc. 2nd Conf. Foodborne Contamination and Egyptian's Health, 23-24 April, El-Mansoura, Egypt.
- Adams, M. R. and M.O. Moss (1995):** Food microbiology., The Royal Society of chemistry.
- Allam, S.M.; A.E. Kinezy; F. El-Tahan and A.M. Soliman (1999-a):** Effect of detoxification of aflatoxin (AFB₁) contaminated rations on blood and carcass characteristics of sheep., Egypt. J. Nutrition and Feed, 2 (Special Issue): 1.
- Allam, S.M.; H.M. El-Hossiény ; A.M. Abdel-Gawad; S.A. El-Saadany and A.M.M. Zeid, (1999-b):** Medicinal herbs and plants as feed additives for ruminants., 1-Effect of using some medicinal herbs and plants as feeds additives on Zaraibi goat performance., Egyptian J. Nutrition and Feeds, 2 (Special Issue): 349.
- Allam, S.M.; A.A. Soliman; M.E. Ahmed and M.A. Shebl (2002):** Aflatoxicosis in Egyptian sheep. 2- clinical chemistry and carcass quality., Proc. 2nd Conf. Foodborne Contamination and Egyptians' Health, 23-24 April, El- Mansoura, Egypt.

- Anonymous, W. S. (1998):** Aflatoxins., WHO Food Additives Series., 40: 359-468.
- Atasever, M.; G. Adiguzel; M. Atasever; H. Özlü and K. Özturan (2010):** Occurrence of Aflatoxin M1 in UHT Milk in Erzurum-Turkey., Kafkas. Univ. Vet. Fak. Derg., 16 (Suppl A): S119-S122.
- Bakirci, I. (2001):** A study on the occurrence of aflatoxin M in milk and milk products produced in Van province of Turkey, Food Control 12 (1), pp. 47–51.
- Barbiroli, A.; F. Bonomi; F. Benedetti; S. Mannino; L. Monti; T. Cattaneo and S. Iametti (2007):** Binding of aflatoxin M1 to different protein fractions in ovine and caprine milk. J. Dairy Sci., 90:532–540.
- Baskaya, R.; A. Aydin; A. Yildiz and K. Bostan (2006):** Aflatoxin M1 levels of some cheese varieties in Turkey., Medycyna Wet., 62: 778–780.
- Bhat, R.V. and J.D. Miller (1991):** Mycotoxins and Food Supply., Food, Nutrition and Agriculture - Food for the Future., FAO.
- Bhat, R.V. and S. Vasanthi (2003):** Food Safety in Food Security and Food Trade: Mycotoxin Food Safety Risk in Developing Countries. Washington D.C. (IFPRI), International Food Policy Research Institute, (Brief 3).
- Biopharm R. and H. Gmb (1999):** Enzyme immunoassay for the quantitative analysis of aflatoxins. M1 Art. No: R-1101 and B1 Art. No: R-1201. R-Biopharm GmbH, Darmstadt, Germany.
- Blunden, G.; O.G. Roch; D.J. Rogers; R.D. Coker; N. Bradburn and A.E. John (1991):** Mycotoxins in food., Med. Lab. Sci., 48 (4): 271-82.
- Brackett, R.E. and E.H. Marth (1982):** Association of aflatoxin M1 with casein. Z Lebensm Unters Forsch, 174: 439-441.
- Brera, A.C.; M. Miraglia and M. Colatosti (1998):** Evaluation of the impact of mycotoxins on human health: Sources of errors., Micro. Chemical Journal, 59 (1): 45-49.
- Cashman, K.D. (2002):** Macrominerals in milk and dairy products, nutritional significance. In: H. Roginski, P.F. Fox and J.W. Fuquay, Editors, Encyclopedia of Dairy Sciences, Academic Press, London, UK, pp. 2051–2058.
- CAST (1989):** Mycotoxins: Economic and Health Risk., Council on Agricultural Science and Technology., Report 116: 1-91.
- Cavaliere, G.; P. Foglia; E. Pastorini; R. Samperi and A. Lagana (2006):** Liquid chromatography/Tandem mass spectrometric confirmatory method for determining aflatoxin M1 in cow milk comparison between electrospray and atmospheric pressure photoionization sources., J. Chromatography A., 1101: 69-78.
- Chen, J. and J. Gao (1993):** The Chinese total diet study in 1990. Part I. Chemical contaminants. J. AOAC Int., 76: 1193- 1205.

- Chifriuc, C.; G. Mihaescu and N.M. Milita (2010):** Aflatoxins health risk factors., *Bacteriol Virusol Parazitol Epidemiol.*, 55 (1):19-24.
- Chu, F.S. (1991):** "Mycotoxins: Food Contamination, Mechanism, Carcinogenic Potential and Preventive Measures." *Mutation Res.*, 259, 291–306.
- Cleveland, T.E.; P.F. Dowd ; A.E. Desjardins; D. Bhatnagar and P.J. Cotty (2003):** United States Department of Agriculture-Agricultural Research Service research on pre-harvest prevention of mycotoxins and mycotoxigenic fungi in US crops., *Pest. Manag. Sci.*, 59: 629–642.
- Creppy, E.E. (2002):** Update of survey, regulation and toxic effects of mycotoxins in Europe. *Toxicology Letters*. 127: 19-28.
- Crosby, D.G. Coordinator (1969):** Symposium on natural food contamination., *J. Agr. Food Chem.*, 17: 413.
- CTA (1997):** Technical Centre for Agricultural and Rural Cooperation Technical., Leaflet No. 3.
- Dutton, M.F.; K. Ehrlich and J.W. Bennett (1985):** Biosynthetic relationship among aflatoxins B1, B2, M1, and M2., *Appl. Environ. Microbiol.*, 49 (6): 1392-5.
- El-Shinawy, S.H.; Abd El-Aziz, A.M. and El-Hady, H.A. (1994):** Microbiological quality of infant powdered Milk., *J. Egypt, Vet. Med. Ass.*, Vol.55, No. 1, 2: 147-154.
- Fallah, A.A. (2010):** Assessment of aflatoxin M1 contamination in pasteurized and UHT milk marketed in central part of Iran. *Food and chemical toxicology an international journal published for the British Industrial Biological Research Association.*, Volume: 48, Issue: 3, Publisher: Elsevier Ltd, Pages: 988-991.
- FAO "Food and Agriculture Organization of the United Nations" (1991):** Food Nutrition and Agriculture – Food for the Future. No., 1. FAO, Rome.
- FAO "Food and Agriculture Organization of the United Nations" (1997-a):** Agriculture food and nutrition for Africa – A resource book for teachers of agriculture. FAO, Rome.
- FAO "Food and Agriculture Organization of the United Nations" (1997-b):** Worldwide regulations for mycotoxins, 1995. A compendium. FAO, Food and Nut. Paper 64, Rome.
- FAO "Food and Agriculture Organization of the United Nations" (1996):** Joint FAO/WHO Expert Committee on Food Additives, 17–26, FAO, Rome.
- FAO "Food and Agriculture Organization of the United Nations" (2001):** Manual on the Application of the HACCP system in Mycotoxin Prevention and Control. FAO, Rome.
- FAO/WHO food standards program (1996):** Joint FAO/WHO Committee Of Government Experts On The Code Of Principles Concerning Milk

And Milk Products., Report of the 22nd Session, CX 5/70., Appendix XV IDF Statement on Aflatoxins in Milk. Rome, Italy.

- Galvano, F.; V. Galofaro and G. Galvano (1996):** Occurrence and stability of aflatoxin M1 in milk and milk products: A worldwide review. *J. Food Protect.*, 59: 1079-1090.
- Galvano, F.; V. Galofaro; A. Ritieni ; M. Bognanno ; A. De Angelis and G. Galvano (1998):** Survey of the occurrence of aflatoxin M1 in dairy products marketed in Italy., *Food J. Prot.* (1), pp. 738–741.
- Galvano, F.; V. Galofaro; A. Ritieni; M. Bognanno; A. De Angelis and G. Galvano, (2001):** Survey of the occurrence of aflatoxin M1 in dairy products marketed in Italy: second year of observation, *Food Additives and Contaminants* 18 (7), pp. 644–646.
- German Cano Sancho; Sonia Marin; Antonio J. Ramos; Juan Peris-Vicente and Vicente Sanchis (2010):** Occurrence of aflatoxin M1 and exposure assessment in Catalonia (Spain)., *Rev Iberoam Micol.*; 27 (3): 130–135., www.elsevier.es/reviberoammicol
- Ghazani, M.H.M. (2009):** Aflatoxin M1 contamination in pasteurized milk in Tabriz (northwest of Iran)., *Food Chem. Toxicol.*, Volume: 47 ISSN: 1873-6351.
- Goto, T.; D.T. Wicklow and Y. Ito (1996):** Aflatoxin and cyclopiazonic acid production by sclerotium producing *Aspergillus tamari* strain., *App. Environ. Microbiol.*, 62: 4036.
- Gulf Standard (1997):** Regulates the maximum limits aflatoxins permitted in foods and animal feeds., No. 841.
- Gundinc, U. and A. Filazi (2009):** Detection of aflatoxin M1 concentrations in UHT milk consumed in Turkey markets by ELISA. *Pak. J. Biol. Sci.*, 12: 653-656.
- Haggag Ola, G.; El Mosallamay A. Waffa; Amin A. Refky and R. Haroun Marcelle (2001):** Induction of apoptosis by aflatoxicosis and its modulation by ascorbic acid and sequential biochemical and histopathological changes in albino rats., *Mansoura J. Forensic Med. Clin. Toxicol.*, 9 (1): 35-52.
- Hollinger, K. and H.E. Ekperigin (1999):** Mycotoxicosis in food producing animals., *Vet. Clin. North. Am. Food Anim. Pract.*, 15: 133–165.
- Hoppe, C.; C. Mølgaard and K.F. Michaelsen (2006):** Cow's Milk and Linear Growth in Industrialized and Developing Countries., *Annual Review of Nutrition* Vol. 26: 131-173.
- Hussein, H.S. and J.M. Brasel (2001):** Toxicity, metabolism, and impact of mycotoxins on humans and animals., *Toxicology.*, 167: 101–134.
- IARC “International Agency for Research on Cancer” (1993):** Monographs on the evaluation of carcinogenic risks to human., Vol.56., Some natural occurring substances: Food items and constituents, heterocyclic aromatic amines and mycotoxins., International Agency for Research on Cancer., Lion.

- IARC “International Agency for Research on Cancer” (2002):** Some Traditional Herbal Medicines, Some Mycotoxins, Naphthalene and Styrene. WHO Press, Geneva, Switzerland. 82: 10-13.
- Ito, Y.; S.W. Peteson; D.T. Wicklow and T. Goto (2001):** *Aspergillus pseudotamarii*, a new aflatoxin-producing species and genetic variation in its sibling species., *A. nomius*., *Mycologia*., 93: 689.
- Jelinek, C.F.; A.E. Pohland and G.E. Wood (1989):** Worldwide occurrence of mycotoxin in foods and feeds, an update., *J. AOAC*, 72: 223-230.
- Kamkar, A. (2008):** Detection of Aflatoxin M1 in UHT milk samples by ELISA., *J. Vet. Res.* 63 (2): 7-12.
- Klick, M.A.; E.J. Mullaney; C.B. Daly and J.W. Cary (2000):** Molecular and physiological aspects of aflatoxin and sterigmatocystin biosynthesis by *Aspergillus tamari* and *A. ochraceoroseus*., *Appl. Microbial. Biotechnol.*, 53: 605.
- Kocabas, C.N. and B.E. Sekerel (2003):** Does systemic exposure to aflatoxin B1 cause allergic sensitization? *Allergy*. 58: 363.
- Kovacs, M. (2004):** Nutritional health aspects of mycotoxins., *Orv. Hetil.*, 145 (34): 1739-46.
- Lafont, P.; M. Siriwardana and J. Lafont (1989):** Genotoxicity of Hydroxyl-aflatoxins M1 and M4. *Micr. Alim. Nut.*, 7: 1-8.
- Lewis, L.; M. Onsongo; H. Njapau; H. Schurz-Rogers; G. Luber; S. Kieszak; J. Nyamongo; L. Backer; A.M. Dahiye and A. Misore (2005):** Aflatoxin contamination of commercial maize products during an outbreak of acute aflatoxicosis in eastern and central Kenya. *Environ. Health Perspect.*, 113: 1763-1767.
- López, C.E.; L.L. Ramos; S.S. Ramadán and L.C. Bulacio (2003):** Presence of aflatoxin M1 in milk for human consumption in Argentina, *Food Control* 14 (1), pp. 31–34.
- Macdonald, S. and I. Castle (1996):** A UK retail survey of aflatoxins in meat and their fate during cooking., *Food. Contam.* 13, 121-128.
- Martins, M.L. and H.M. Martins (2000):** Aflatoxin M1 in raw and ultra-high temperature-treated milk commercialized in Portugal, *Food Additives and Contaminants* 17 (10), pp. 871–874.
- Montesano, R.; P. Hainaut and C.P. Wild (1997):** Hepatocellular carcinoma: From gene to public health., Review, *Journal of National Cancer Institute*., 89: 1844-51.
- Moss, M.O. (1998):** Recent studies of mycotoxins, *Journal of Applied Microbiology, Symposium Supplement* 84, pp. 62–76.
- Moustafa, R.M. (1994):** Mycotoxin residues in meat and meat products., Ph. D. Thesis, Fac. Vet .Med., Cairo Univ., Egypt.

- Neal, G.E.; D.L. Eaton; D.J. Judah and A. Verna (1998):** Metabolism and Toxicity of aflatoxins M1 and B1 in human-derived in vitro systems. *Toxicol. Appl. Pharm.*, 151 (1): 152-158.
- Oliveira, C.A.F. and J.C.O. Ferraz (2007):** Occurrence of aflatoxin in pasteurised, UHT milk and milk powder from goat origin, *Food Control* 18 (4), pp. 375–378.
- Orriss, G.D. (1997):** Animal diseases of public health importance., *Emerg. Inf. Diseases.*, 3: 497–502.
- Park, D.L. (2002):** Effect of processing on aflatoxin. *Adv. Exp. Med. Biol.*, 504: 173-179.
- Peraica, M.; B. Radic; A. Lucic and J. Pavlovic (1999):** World health organ., 77-754.
- Peterson, S.W.; Y. Ito ; B.W. Horn and T. Goto (2001):** *Aspergillus bombycis*, a new aflatoxigenic species in *Aspergillus* section *flavi*., *Mycol. Res.*, 105: 233.
- Pipet, A. (1998):** Natural Occurrence of Mycotoxins in Foods and Feeds: An Update products. *Review. Rev. de Med. Veter.*, 6: 479-492.
- Pitt, J.I. (2000):** Toxigenic fungi: which are important., *Med. Mycol.*, 38 Suppl., 1: 17-22.
- Prandini, A; G. Tansini; S. Sigolo; L. Filippi; M. Laporta and G. Piva (2009):** On the occurrence of aflatoxin M1 in milk and dairy products. *Food Chem Toxicol.* May; 47 (5): 984-91.
- Qian, G.S.;P. Yasei and G.C. Young (1984):** “Rapid Extraction and Detection of AFM1 in Cow's Milk by HPLC and RIA.” *Anal. Chem.*, 56: 2079–2080.
- Rahimi, E. and G. Karim (2008):** Determination of aflatoxin M1 in raw milk producing in dairy farmsin Chahar-Mahal-Bakhtiari province using ELISA., *J. Food Sci. Technol.*, 5: 51–59.
- Roussi, V.; A. Govaris; A. Varagouli and N.A. Botsoglou. (2002):** Occurrence of aflatoxin M1 in raw and market milk commercialized in Greece. *Food Addit Contam. Sep*;19 (9): pp. 863-8.
- SASO "Saudi Arabia Standard Organization" (1996):** 1151/1996 Maximum limits of Mycotoxins permitted in foods and animal feeds-Alfa-toxins.
- Sassahara, M.; D.P. Netto and E.K. Yanaka (2005):** Aflatoxin occurrence in foodstuff supplied to dairy cattle and aflatoxin M1 in raw milk in the North of Parana state. *Food Chem. Toxicol.*, 43: 981-984.
- Sayed, M.A.; E.A. Mohamoud and A.A. Abou El-Alla (2000):** Mycoflora and natural occurrence of mycotoxin in meat of imported bulls, poultry and some meat products., *Assuit Vet. Med. J.*, 43 (86): 188-200.
- Sherif, S.O. (2003):** Aflatoxins and child health: towards a strategy for prevention and control., *Med. J. Cairo Univ.*, 71 (2 Suppl., 2): 1-14.

- Smith, J.E.; C.W. Lewis; J.G. Anderson and G.L. Solomon (1994):** Mycotoxins in human nutrition and health., European Commission DGXII, Science Research and Development, EUR 16048 EN.
- SPSS, (1997):** Statistical Package for Social Science. SPSS®, for windows release 8.0 SPSS (Inc.), 444N, Michigan Avenue, Chicago, IL, 60611, USA.
- Stoloff, L. (1977):** Aflatoxins: An overview. In: Proceeding of a conference on mycotoxins in human and animal health., pp. 7-28. Park Forest South: Pathotox Puplisher Inc.
- Takahashi-Ando, M.; S. Ohsato; T. Shibata; H. Hamamoto; I. Yamaguchi and M. Kimura (2004):** Metabolism of zearalenone genetically modified organism expressing the detoxification gene from *clonostachys tosea.*, *Appl. Environ. Microbio.*, 70 (6): 3239-3245.
- Tekinşen, K.K. and H.S. Eken (2008):** Aflatoxin M1 levels in UHT milk and kashar cheese consumed in Turkey., *Food and Chemical Toxicology.*, Volume 46, Issue 10, October, Pages 3287-3289.
- Trucksess, M.; C. Weaver; C. Oles; K. D'Ovidio and J. Rader (2006):** Determination of aflatoxins and ochratoxin A in ginseng and other botanical roots by immunoaffinity column cleanup and liquid chromatography with fluorescence detection., *J. AOAC Int.*, 89 (3): 624-30.
- Unusan, N. (2006):** Occurrence of aflatoxin M1 in UHT milk in Turkey, *Food and Chemical Toxicology.*, 44 (11), pp. 1897–1900.
- USFAD "United State Food and Drug Administration" (2000):** Foodborne Pathogenic Microorganisms and Natural Toxins Handbook. <http://vm.cfsan.fda.gov/~mow/chap41.html>.
- Van Egmond, H.P. (1989):** “Aflatoxin M1: Occurrence, Toxicity, Regulation”, *Mycotoxins in Dairy Products*, Van Egmond, H.P. (Ed.). London: Elsevier Applied Science Publishers Ltd., 11-49.
- Van Egmond, H.P. (1995):** Current situation on regulations for mycotoxins. Overview of tolerances and status of standard method of sampling and analysis. *Food Addit. Contam.*, 6: 139-188.
- Van Egmond, H.P.; U.K. Svensson and J.M. Fremy (1997):** Mycotoxins. In: *Residues and contaminants in milk and milk products (special Issue 9701)*, Brussels, International Dairy Federation., pp. 17-88.
- Varman, A.H. and M.G. Evans (1991):** *Food borne pathogens.*, [an illustrated textbook] 2nd Ed., Wolfe Publishing Ltd., England.
- WHO "World Health Organization" (1999):** *Basic Food Safety for Health Workers.*
- WHO "World Health Organization" (2002):** *Evaluation of Certain Mycotoxins in Food.* World Health Organization, Technical Report Series No. 906, WHO, Geneva, Switzerland., pp. 1-62.

Yaroglu, T. ; H.H. Oruc and M. Tayar (2005): Aflatoxin M1 levels in cheese samples from some provinces of Turkey, Food Control 16: (10), pp. 883–885.

تقدير تركيزات الأفلاتوكسين إم ١ في الحليب البقري كامل الدسم المُعامل بالحرارة الفائقة المُباع للاستهلاك في نجران-السعودية مع بيان الأهمية الصحية

مدحت إبراهيم محمد عبدالله^١، محمد صالح بزالو^٢، محمد بن زيد ناصر الجليفي^٣

تمت هذه الدراسة بدعم من عمادة البحث العلمي بجامعة نجران (مسجل برقم: NU/١١/٦٧)

^١ أستاذ مشارك ميكروبيولوجي في برنامج تقنية المختبرات الطبية، قسم العلوم الطبية التطبيقية، كلية المجتمع، جامعة نجران، نجران، المملكة العربية السعودية.

^٢ أخصائي أول في الطب البيطري وباحث في فحوص صحة الأغذية بمختبر ميناء دمياط البحري، المختبر المرجعي لتحليل سلامة الأغذية من أصل حيواني، معهد بحوث صحة الحيوان، مركز البحوث الزراعية، الجيزة، مصر، وعضو بمختبر السموم والهرمونات ويقايا الأدوية، وزارة الزراعة السعودية، الرياض.

^٣ أستاذ العلوم الجزيئية والتكنولوجيا الحيوية ومدير المشروع الوطني (UTF/SAU/035/SAU) لتحسين الأمراض الحيوانية وإنتاج اللقاح في المملكة العربية السعودية بالتعاون مع منظمة الأغذية والزراعة، والمدير العام لمختبرات الطب البيطري، وزارة الزراعة السعودية، الرياض.

أجريت هذه الدراسة للكشف عن تواجد مادة الأفلاتوكسين إم ١ السامة في الحليب البقري كامل الدسم المُعقم بالحرارة الفائقة (UHT) المُسوق للاستهلاك في مدينة نجران بالسعودية، وذلك باستخدام تقنية الإليزا ELISA (فحص إنزيمات المناعة المرتبطة)، وقد تم إجراء الدراسة في الفترة من سبتمبر ٢٠١١م إلى يناير ٢٠١٢م على عدد ٩٦ عينة عشوائية من حليب الإبقار كاملة الدسم المعامل بالحرارة الفائقة بـ UHT، حيث دلت النتائج على تواجد سُم AFM1 في ٧٩ عينة (٨٢,٣٠%) بتركيزات تتراوح بين 0.01-0.19 ميكروجرام لكل لتر، وبمتوسط قدرة 0.058 ± 0.0053 ميكروجرام لكل لتر، ومن ناحية أخرى تبين أن مستويات التلوث بـ AFM1 في عينات الحليب المعقم (UHT) منخفضة ولا تتجاوز الحد الأقصى المسموح به من هيئة النقييس الموحد السعودية، وبالتالي لا تشكل خطورة على صحة المستهلك في مدينة نجران بالسعودية، هذا وقد تم مناقشة النتائج والأهمية الصحية ومدى خطورة وتأثير سُم الأفلاتوكسين إم ١ على صحة الإنسان وما يجب أن يُتبع من إجراءات للمساهمة في إنتاج أقل تلوثاً بالسموم الفطرية.

^١ باحث أول في فحوص صحة الأغذية بمختبر ميناء دمياط البحري، المختبر المرجعي لتحليل سلامة الأغذية من أصل حيواني، معهد بحوث صحة الحيوان، مركز البحوث الزراعية، الجيزة، مصر.